

Scaling and asymptotic scaling in two-dimensional CP^{N-1} models *

Massimo Campostrini, Paolo Rossi, and Ettore Vicari

*Istituto Nazionale di Fisica Nucleare, Sezione di Pisa,
Dipartimento di Fisica dell'Università, I-56126 Pisa, Italy.*

October 6th, 1992

Abstract

Two-dimensional CP^{N-1} models are investigated by Monte Carlo methods on the lattice, for values of N ranging from 2 to 21. Scaling and rotation invariance are studied by comparing different definitions of correlation length ξ . Several lattice formulations are compared and shown to enjoy scaling for ξ as small as 2.5.

Asymptotic scaling is investigated using as bare coupling constant both the usual β and β_E (related to the internal energy); the latter is shown to improve asymptotic scaling properties. Studies of finite size effects show their N -dependence to be highly non-trivial, due to the increasing radius of the $\bar{z}z$ bound states at large N .

1 Introduction

Two-dimensional CP^{N-1} models share many dynamical features with four-dimensional $\text{SU}(N)$ gauge theories, but are much easier to explore numerically and are amenable to analytical analysis by means of the $1/N$ expansion. CP^{N-1} models can therefore be useful as a test-bed for algorithms and numerical methods to be applied to lattice QCD, since systematic errors can be explored in details and numerical results can be compared with precise theoretical predictions. Indeed several studies of CP^{N-1} and related models have been presented in this Conference.

In this paper we summarize studies of the scaling, asymptotic scaling, and finite-size scaling properties of different lattice formulations of CP^{N-1} models.

*Talk presented at the *Lattice '92* Conference, Amsterdam.

The full details of the simulations and analyses of different aspects of the models can be found in Refs. [1, 2].

Many recent lattice studies of CP^{N-1} models used the “quartic” action

$$S_1 = -N\beta \sum_{n,\mu} |\bar{z}_{n+\mu} z_n|^2. \quad (1)$$

We performed most of our simulations using the “gauge” action

$$S_g = -N\beta \sum_{n,\mu} (\bar{z}_{n+\mu} \lambda_{n,\mu} z_n + \bar{z}_n \bar{\lambda}_{n,\mu} z_{n+\mu}). \quad (2)$$

z_n is an N -component complex scalar field constrained by the condition $\bar{z}_n z_n = 1$, and $\lambda_{n,\mu}$ is a $\text{U}(1)$ gauge field. S_g enjoys several advantages over S_1 , the most important being a better scaling behavior. We also considered the tree-level Symanzik-improved versions of both actions, which will be denoted by S_1^{Sym} and S_g^{Sym} .

We performed Monte Carlo simulations of the CP^1 , CP^3 , CP^9 , and CP^{20} models, for values of the correlation length ξ ranging from 2.5 to 30.

2 Correlation Length and Scaling

The most interesting correlation function of the model selects the $\text{SU}(N)$ -adjoint, gauge-neutral channel:

$$\begin{aligned} G_P(x) &= \langle \text{Tr}\{P(x)P(0)\} \rangle_{\text{conn}}, \\ P_{ij}(x) &= \bar{z}_i(x) z_j(x). \end{aligned} \quad (3)$$

We defined the usual wall-wall correlation length ξ_w from the exponential decay of the zero-momentum projection of G_P . In order to check for rotation invariance, we also defined the diagonal wall-wall correlation length ξ_{dw} in the obvious way. Both ξ_w and ξ_{dw} tend to the inverse mass gap in the scaling limit. The ratio ξ_{dw}/ξ_w is plotted in Figs. 1a, 2a, 3, and 4 for different values of N and different actions, showing rotation invariance even for the smallest values of ξ .

We also chose a different definition of correlation length:

$$\xi_G^2 = \frac{1}{4 \sin^2 \pi/L} \left[\frac{\tilde{G}_P(0,0)}{\tilde{G}_P(0,1)} - 1 \right], \quad (4)$$

where $\tilde{G}_P(p_1, p_2)$ is the Fourier transform of G_P ; $\tilde{G}_P(0,0)$ is the magnetic susceptibility χ . In the scaling limit, ξ_G tends to the second moment of the correlation function. The physical quantity ξ_G/ξ_w must be independent of β and on the choice of the action in the scaling region. ξ_G/ξ_w is plotted in Figs. 1b, 2b, 3, and 4 for different values of N and different actions, showing scaling for all the values of ξ . In the following we will choose ξ_G as length scale, since it is less noisy than ξ_w or ξ_{dw} .

3 Asymptotic Scaling

We compared the β -dependence of the lattice mass scale $1/\xi_G$ with the two-loop perturbative formula $f(\beta) = (2\pi\beta)^{2/N} \exp(-2\pi\beta)$, by considering the ratio $M_G/\Lambda_{\text{latt}} \equiv [\xi_G f(\beta)]^{-1}$. In the asymptotic regime, $M_G/\Lambda_{\text{latt}}$ must be constant plus corrections $O(1/\beta)$; the $\beta \rightarrow \infty$ limit may therefore be approached very slowly. In order to compare different lattice formulations, we reexpressed the specific Λ_{latt} of each formulation in terms Λ_g .

We used two different schemes for the analysis: in the “standard” scheme the inverse bare coupling is simply β , the action parameter. In the “energy” scheme, suggested by G. Parisi [3], the inverse bare coupling is $\beta_E \equiv 1/(2E)$; E is the expectation value of the link action, normalized to $E(\beta = \infty) = 0$. At the perturbative level, β and β_E are simply two different definitions of coupling constant. We reexpressed Λ_{latt} in the “energy” scheme in terms of Λ_g in the “standard” scheme.

The results for CP^1 , CP^3 , and CP^9 , are presented in Figs. 5–7. Results for the “standard” scheme show asymptotic scaling violations and disagreement between different lattice actions, but both effects are compatible with the expected $1/\beta$ behavior. On the other hand, results for the “energy” scheme are remarkably β - and action-independent. In the case of CP^1 , the “energy” scheme result is in agreement with the exact result of Ref. [4]:

$$\frac{M_G}{\Lambda_g} = \frac{8}{e} \sqrt{32} \exp\left(\frac{\pi}{4}\right) \cong 36.51. \quad (5)$$

4 Finite Size Scaling

We examined finite size scaling effects by studying the finite size scaling function of an observable $\mathcal{O}$: $f_{\mathcal{O}}(L/\xi) \equiv \mathcal{O}_L(\beta)/\mathcal{O}_{\infty}(\beta)$, where $\mathcal{O}_L$ is the expectation value of the observable $\mathcal{O}$ measured on a $L \times L$ lattice. In the scaling region, $f_{\mathcal{O}}$ must be a *universal* function of L/ξ , independent of β and of the lattice action. We present here results for the magnetic susceptibility χ ; f_{ξ} has a behavior very similar to f_{χ} .

We expect a qualitative difference in the finite size behavior of CP^{N-1} models at small and large N . Finite size effects for CP^1 are expected to be dominated by the inverse mass gap, therefore following the predictions of Ref. [5]: $f_{\mathcal{O}} = 1 + O(\exp\{-L/\xi\})$. f_{χ} is plotted in Fig. 8, and it is in agreement with the above prediction.

No clear theoretical predictions are available for the CP^3 model. f_{χ} is plotted in Fig. 9.

The large- N expansion predicts a radius of the lowest-lying state proportional to $\xi N^{1/3}$ [6]. Therefore we expect, for N large enough, finite size effects to be dominated by the bound state radius rather than by the inverse mass gap. Quantities like ξ and χ are expected to increase when L is decreased, in contrast

with the CP^1 case. Moreover the function $f_{\mathcal{O}}(L/(\xi N^{1/3}))$ should not depend on N (at least for $N \lesssim 100$; for huge values of N , the bound state decouples from χ and ξ_G). Fig. 10 shows f_{χ} plotted as a function of $L/(\xi N^{1/3})$, for both CP^9 and CP^{20} , and it is in agreement with the large- N expansion.

References

- [1] M. Campostrini, P. Rossi, and E. Vicari, Phys. Rev. **D46** (1992) 2647.
- [2] M. Campostrini, P. Rossi, and E. Vicari, Pisa preprint IFUP-TH 34/92, Phys. Rev. **D**, in press.
- [3] G. Parisi, in *High Energy Physics—1980*, Proceedings of the XXth Conference, Madison, Wisconsin, 1980, AIP Conf. Proc. No. 68 (AIP, New York, 1981).
- [4] P. Hasenfratz, M. Maggiore, and F. Niedermayer, Phys. Lett. **B245** (1990) 522.
- [5] M. Lüscher, Phys. Lett. **118B** (1982) 391.
- [6] E. Witten, Nucl. Phys. **B149** (1979) 285.

Figure captions

Fig. 1: Scaling tests for CP^1 .

Fig. 2: Scaling tests for CP^3 .

Fig. 3: Scaling tests for CP^9 .

Fig. 4: Scaling tests for CP^{20} .

Fig. 5: Asymptotic scaling tests for CP^1 .

Fig. 6: Asymptotic scaling tests for CP^3 .

Fig. 7: Asymptotic scaling tests for CP^9 .

Fig. 8: Finite size scaling for CP^1 .

Fig. 9: Finite size scaling for CP^3 .

Fig. 10: Finite size scaling for CP^9 and CP^{20} .